

### (15) Mortalidad por cirrosis hepática <sup>1/</sup>

Entidad Federativa	Hombres	Mujeres	Total
<b>Nacional</b>	<b>33,47</b>	<b>10,22</b>	<b>21,79</b>
Aguascalientes	20,93	4,75	12,64
Baja California	27,51	5,96	16,91
Baja California Sur	18,30	4,20	11,44
Campeche	34,14	14,08	24,25
Coahuila	21,98	7,54	14,81
Colima	34,21	6,85	20,55
Chiapas	30,07	12,70	21,50
Chihuahua	24,96	5,94	15,56
Distrito Federal	30,76	10,05	20,15
Durango	16,09	3,98	10,00
Guanajuato	26,39	7,83	16,87
Guerrero	24,72	5,99	15,29
Hidalgo	61,25	17,76	39,14
Jalisco	24,90	8,62	16,70
México	34,98	11,10	23,03
Michoacán	29,79	7,57	18,44
Morelos	39,61	12,05	25,73
Nayarit	17,98	7,14	12,61
Nuevo León	19,92	8,24	14,12
Oaxaca	49,13	12,19	30,38
Puebla	62,86	18,12	40,19
Querétaro	42,64	11,87	27,15
Quintana Roo	26,11	8,25	17,57
San Luis Potosí	29,16	7,57	18,22
Sinaloa	16,02	4,35	10,26
Sonora	21,24	4,98	13,23
Tabasco	27,14	5,80	16,52
Tamaulipas	21,56	7,26	14,45
Tlaxcala	39,12	16,74	27,85
Veracruz	57,23	15,39	36,12
Yucatán	50,97	28,31	39,62
Zacatecas	18,82	4,28	11,38

<sup>1/</sup> Mortalidad observada por cada 100,000 habitantes.

Fuentes: Base de datos de las defunciones INEGI/SSA, 2004 y Proyecciones de Población 2000-2050 del CONAPO, 2002.