

(15) Mortalidad por cirrosis hepática ^{1/}

Entidad Federativa	Hombres	Mujeres	Total
Nacional	32,78	10,54	21,60
Aguascalientes	19,63	6,17	12,74
Baja California	26,02	5,87	16,11
Baja California Sur	22,46	7,37	15,12
Campeche	31,80	13,59	22,82
Coahuila	23,73	9,35	16,59
Colima	35,83	7,11	21,48
Chiapas	30,18	12,74	21,51
Chihuahua	24,18	6,43	15,41
Distrito Federal	30,28	10,24	20,00
Durango	19,92	8,44	14,15
Guanajuato	28,92	7,15	17,75
Guerrero	25,33	7,80	16,50
Hidalgo	56,17	17,53	36,53
Jalisco	26,41	8,63	17,46
México	31,83	10,24	21,03
Michoacán	29,52	8,01	18,52
Morelos	37,44	16,41	26,85
Nayarit	17,86	5,87	11,93
Nuevo León	21,02	7,97	14,54
Oaxaca	46,53	13,81	29,89
Puebla	63,04	19,25	40,85
Querétaro	40,88	10,54	25,61
Quintana Roo	23,41	8,53	16,22
San Luis Potosí	28,62	7,37	17,85
Sinaloa	16,03	4,46	10,32
Sonora	18,30	7,27	12,87
Tabasco	27,14	9,22	18,22
Tamaulipas	21,87	7,82	14,89
Tlaxcala	38,88	15,56	27,14
Veracruz	55,40	15,06	35,05
Yucatán	51,43	26,29	38,84
Zacatecas	17,36	6,06	11,57

^{1/} Mortalidad observada por cada 100,000 habitantes.

Fuentes: Cifras preliminares del SEED 2005; DGIS, SSA, y Proyecciones de Población 2000-2050 del CONAPO, 2002.