

(6) Mortalidad por enfermedades diarreicas en menos de 5 años ^{1/}

Entidad federativa	Hombres	Mujeres	Total
E. U. M	19.86	17.49	18.70
Aguascalientes	1.86	9.65	5.67
Baja California	8.44	2.96	5.76
Baja California Sur	16.91	4.48	10.88
Campeche	12.91	2.70	7.93
Coahuila	10.85	5.25	8.12
Colima	15.27	7.94	11.67
Chiapas	53.55	49.60	51.62
Chihuahua	18.40	12.24	15.40
Distrito Federal	12.48	13.24	12.85
Durango	9.24	5.50	7.41
Guanajuato	19.35	17.81	18.60
Guerrero	27.54	13.19	20.53
Hidalgo	11.41	8.25	9.87
Jalisco	11.72	9.03	10.40
México	22.92	22.26	22.60
Michoacán	14.71	15.37	15.03
Morelos	21.53	15.84	18.74
Nayarit	10.71	13.45	12.05
Nuevo León	5.69	4.87	5.29
Oaxaca	41.19	35.17	38.24
Puebla	38.25	32.24	35.31
Queretaro	15.32	15.96	15.63
Quintana Roo	12.64	18.92	15.71
San Luis Potosí	18.37	20.12	19.22
Sinaloa	3.90	4.90	4.39
Sonora	9.34	9.80	9.56
Tabasco	23.90	14.56	19.33
Tamaulipas	6.12	5.69	5.91
Tlaxcala	30.97	22.14	26.64
Veracruz	19.96	22.45	21.18
Yucatán	28.81	21.35	25.16
Zacatecas	15.60	16.22	15.90

1/ Tasa de mortalidad observada por 100,000 habitantes de ese grupo de edad. Fuente: Base de datos de las defunciones INEGI/SS y SEED 2005, Dirección General de Información en Salud, SS. Proyecciones de la Población 2000-2050 del CONAPO, 2002.