

(12a) Mortalidad por enfermedades isquémicas del corazón, observada ^{1/}

Entidad federativa	Hombres	Mujeres	Total
E. U. M	56.07	43.92	49.96
Aguascalientes	40.05	36.10	38.03
Baja California	64.84	44.31	54.75
Baja California Sur	69.31	39.28	54.71
Campeche	60.80	33.71	47.44
Coahuila	83.37	59.42	71.49
Colima	66.93	45.00	55.97
Chiapas	29.10	22.44	25.79
Chihuahua	83.06	59.11	71.23
Distrito Federal	81.93	78.44	80.14
Durango	70.76	47.83	59.23
Guanajuato	51.18	39.81	45.37
Guerrero	30.89	26.86	28.86
Hidalgo	55.24	43.38	49.21
Jalisco	60.54	48.39	54.44
México	36.54	30.68	33.61
Michoacán	48.34	36.57	42.32
Morelos	46.01	33.98	39.95
Nayarit	68.08	52.25	60.24
Nuevo León	82.26	54.11	68.28
Oaxaca	39.53	32.38	35.89
Puebla	37.23	31.94	34.55
Queretaro	48.43	37.09	42.72
Quintana Roo	24.29	15.54	20.06
San Luis Potosí	55.04	50.03	52.50
Sinaloa	71.17	50.09	60.77
Sonora	91.09	56.59	74.10
Tabasco	42.63	29.60	36.14
Tamaulipas	74.65	49.99	62.39
Tlaxcala	30.62	21.85	26.21
Veracruz	58.61	46.61	52.55
Yucatán	74.92	65.83	70.37
Zacatecas	55.70	48.23	51.88

1/ Mortalidad observada por cada 100,000 habitantes. Fuente: Base de datos de las defunciones INEGI/SS y SEED 2005, Dirección General de Información en Salud, SS. Proyecciones de la Población 2000-2050 del CONAPO, 2002.