

(15) Mortalidad por cirrosis hepática, observada ^{1/}

Entidad federativa	Hombres	Mujeres	Total
E. U. M	33.47	10.65	22.00
Aguascalientes	17.67	6.36	11.88
Baja California	23.95	5.66	14.96
Baja California Sur	23.62	9.00	16.51
Campeche	31.55	15.16	23.46
Coahuila	25.92	9.67	17.85
Colima	33.47	6.77	20.12
Chiapas	33.33	13.47	23.52
Chihuahua	25.27	6.25	15.88
Distrito Federal	29.19	9.91	19.30
Durango	20.31	7.16	13.70
Guanajuato	28.92	7.69	18.04
Guerrero	23.60	7.98	15.73
Hidalgo	53.11	18.85	35.69
Jalisco	26.89	8.10	17.43
México	35.68	11.10	23.38
Michoacán	25.11	6.85	15.78
Morelos	35.68	15.49	25.51
Nayarit	19.25	6.48	12.93
Nuevo León	21.12	7.83	14.52
Oaxaca	51.13	14.34	32.42
Puebla	64.69	18.89	41.48
Queretaro	41.39	11.04	26.11
Quintana Roo	25.89	8.53	17.50
San Luis Potosí	25.42	8.52	16.85
Sinaloa	15.53	4.02	9.85
Sonora	18.22	7.02	12.71
Tabasco	25.40	8.25	16.86
Tamaulipas	20.48	7.63	14.10
Tlaxcala	40.20	15.56	27.79
Veracruz	59.52	15.60	37.34
Yucatán	51.65	26.40	39.00
Zacatecas	14.90	5.65	10.16

1/ Mortalidad observada por cada 100,000 habitantes. Fuente: Base de datos de las defunciones INEGI/SS y SEED 2005, Dirección General de Información en Salud, SS. Proyecciones de la Población 2000-2050 del CONAPO, 2002.