

### (15) Mortalidad por cirrosis hepática a/

Entidad Federativa	Hombres	Mujeres	Total
<b>Nacional</b>	<b>33.50</b>	<b>10.26</b>	<b>21.84</b>
Aguascalientes	19.36	3.73	11.35
Baja California	24.74	6.23	15.63
Baja California Sur	12.21	6.63	9.49
Campeche	26.52	17.06	21.85
Coahuila	23.15	7.25	15.26
Colima	26.30	7.04	16.86
Chiapas	30.09	11.47	20.84
Chihuahua	22.02	5.54	13.90
Distrito Federal	30.18	10.41	20.07
Durango	18.43	6.23	12.31
Guanajuato	30.40	7.06	18.43
Guerrero	23.66	6.30	14.93
Hidalgo	57.59	17.23	37.08
Jalisco	26.45	8.63	17.50
México	38.88	13.06	25.99
Michoacán	26.72	7.64	17.00
Morelos	36.84	13.20	25.02
Nayarit	19.65	6.20	13.10
Nuevo León	19.98	7.22	13.64
Oaxaca	53.66	13.72	33.45
Puebla	63.14	18.26	40.43
Querétaro	41.06	9.31	25.09
Quintana Roo	22.83	6.56	15.16
San Luis Potosí	24.94	5.91	15.30
Sinaloa	12.94	4.81	8.94
Sonora	20.11	6.34	13.33
Tabasco	23.69	6.56	15.18
Tamaulipas	20.61	6.19	13.48
Tlaxcala	38.27	12.63	25.35
Veracruz	57.27	13.46	35.20
Yucatán	46.36	28.17	37.25
Zacatecas	18.87	3.74	11.13

a/ Tasa de mortalidad observada por 100 000 habitantes de ese grupo de edad. Fuente: Base de mortalidad INEGI-SSA, 2002 y proyecciones de población 2000-2050 del CONAPO, 2002.