

(6) Mortalidad por enfermedades diarreicas en menores de 5 años a/

Entidad Federativa	Hombres	Mujeres	Total
Nacional	22.03	19.22	20.65
Aguascalientes	12.26	12.76	12.50
Baja California	13.76	5.05	9.51
Baja California Sur	4.19	13.28	8.61
Campeche	17.20	2.57	10.05
Coahuila	14.24	8.28	11.33
Colima	11.06	7.67	9.40
Chiapas	42.86	57.06	49.82
Chihuahua	12.34	10.47	11.43
Distrito Federal	13.03	10.36	11.72
Durango	3.62	3.77	3.69
Guanajuato	23.69	15.15	19.51
Guerrero	20.68	18.35	19.54
Hidalgo	20.57	20.60	20.58
Jalisco	12.37	13.52	12.93
México	29.14	27.00	28.09
Michoacán	16.53	13.05	14.83
Morelos	22.41	15.99	19.27
Nayarit	20.23	16.92	18.61
Nuevo León	6.46	4.16	5.34
Oaxaca	48.23	40.96	44.67
Puebla	49.72	33.12	41.59
Querétaro	20.79	16.56	18.72
Quintana Roo	14.64	7.66	11.23
San Luis Potosí	20.03	20.12	20.07
Sinaloa	3.64	9.14	6.33
Sonora	9.76	5.96	7.91
Tabasco	20.95	20.87	20.91
Tamaulipas	9.91	11.05	10.47
Tlaxcala	35.55	21.41	28.62
Veracruz	23.26	20.14	21.73
Yucatán	30.36	24.40	27.45
Zacatecas	11.66	10.79	11.24

a/ Tasa de mortalidad observada por 100 000 habitantes de ese grupo de edad.

Fuente: Base de mortalidad INEGI-SSA, 2002 y proyecciones de población 2000-2050 del CONAPO, 2002.