

(15) Mortalidad por cirrosis hepática b/

Entidad Federativa	Hombres	Mujeres	Total
Nacional	34.03	10.46	22.19
Aguascalientes	20.26	4.63	12.25
Baja California	27.40	6.64	17.19
Baja California Sur	24.14	7.32	15.94
Campeche	28.68	14.86	21.86
Coahuila	24.68	8.62	16.71
Colima	30.82	7.99	19.42
Chiapas	31.14	12.04	21.65
Chihuahua	28.11	5.25	16.81
Distrito Federal	32.71	10.00	21.08
Durango	18.48	5.68	12.05
Guanajuato	29.98	7.19	18.29
Guerrero	23.37	7.37	15.33
Hidalgo	50.86	16.83	33.56
Jalisco	25.83	8.52	17.12
México	39.34	12.95	26.14
Michoacán	27.55	7.88	17.51
Morelos	37.28	13.15	25.14
Nayarit	21.32	6.57	14.02
Nuevo León	16.27	8.52	12.43
Oaxaca	47.46	10.35	28.59
Puebla	63.23	18.64	40.64
Querétaro	45.25	11.58	28.30
Quintana Roo	16.61	8.15	12.52
San Luis Potosí	27.34	7.61	17.35
Sinaloa	14.06	5.51	9.84
Sonora	24.03	5.98	15.15
Tabasco	28.43	6.37	17.47
Tamaulipas	20.41	7.66	14.07
Tlaxcala	38.02	13.76	25.80
Veracruz	57.98	14.47	36.03
Yucatán	52.79	27.44	40.10
Zacatecas	13.05	4.29	8.56

b/ Tasa de mortalidad observada por 100,000 habitantes.

Fuente: Base de mortalidad INEGI-SSA, 2003 y proyecciones de población 2000-2050 del CONAPO 2002.