

**(6) Mortalidad por enfermedades
diarreicas en menores de 5 años c/**

Entidad Federativa	Hombres	Mujeres	Total
Nacional	30.71	24.96	27.90
Aguascalientes	17.88	20.46	19.15
Baja California	24.27	14.56	19.54
Baja California Sur	8.42	40.09	23.81
Campeche	17.51	13.09	15.35
Coahuila	12.89	7.60	10.31
Colima	7.50	15.60	11.47
Chiapas	73.31	65.35	69.41
Chihuahua	18.51	21.30	19.87
Distrito Federal	22.62	16.13	19.44
Durango	8.70	7.77	8.25
Guanajuato	30.94	21.31	26.23
Guerrero	21.45	22.41	21.92
Hidalgo	13.50	9.68	11.63
Jalisco	22.39	17.55	20.02
México	43.41	36.57	40.06
Michoacán	22.63	19.76	21.23
Morelos	24.61	20.52	22.61
Nayarit	26.85	12.97	20.07
Nuevo León	18.61	12.63	15.69
Oaxaca	56.25	43.75	50.13
Puebla	48.13	37.65	43.00
Querétaro	24.89	22.04	23.49
Quintana Roo	23.78	28.74	26.20
San Luis Potosí	28.12	17.63	22.99
Sinaloa	11.18	10.92	11.06
Sonora	19.81	25.95	22.80
Tabasco	42.73	24.34	33.73
Tamaulipas	6.67	7.68	7.16
Tlaxcala	51.35	31.64	41.69
Veracruz	33.03	27.16	30.16
Yucatán	41.42	24.75	33.27
Zacatecas	18.71	8.34	13.63

c/ Tasa de mortalidad observada por 100 000 habitantes de ese grupo de edad.

Fuente: Base de mortalidad INEGI-SSA, 2003 y proyecciones de población 2000-2050 del CONAPO, 2002.